

Name: \_\_\_\_\_

Quest for Quantum

1. In your answers to the following questions, be sure to **INCLUDE THE NAME OF THE PRINCIPLE** that you have used as a part of your answers:
  - a) When filling orbitals around an atom that happen to be of equal energy (i.e. the values of  $n$  and  $l$  are the same) what happens as you add electrons to fill the orbitals?
  
  - b) When adding electrons to a bare nucleus to build an atom, what orbital would fill first?
  
  - c) If you knew how much energy was required to remove an electron from the surface of a metal (Electron Binding Energy) and you knew the exact energy (Photon Energy) of the type of light photon that you were using to perform the removal, how could you calculate the kinetic energy (Electron Kinetic Energy) of the electron? Show a simple equation for your answer in addition to the principle at work.
  
  - d) Why is it difficult to know both the location and the momentum of an electron through any possible experimental technique?

- |                                    |                                      |
|------------------------------------|--------------------------------------|
| - absorption spectra               | - Hund's Rule                        |
| - Aufbau principle                 | - law of conservation of mass energy |
| - atomic absorption                | - line spectra matter waves          |
| - Bohr Model                       | - Pauli Exclusion Principle          |
| - cathode ray                      | - photoelectric effect               |
| - de Broglie wavelength            | - quantum hypothesis                 |
| - emission spectra                 | - Rutherford Model                   |
| - Heisenberg uncertainty principle | - spin coupling                      |

2. Do you know your quantum numbers? List in the order that the quantum numbers were presented in class.

Symbol	Allowed Values (Use Set Notation)	Physical Properties And/or Name

3. Write the complete electron configuration of Caldwellium, element number 111 (also known as ununium).

4. Complete the following table by filling in whatever is missing:

element	n	l	$m_l$	$m_s$	end of config.
${}_{16}\text{S}$					
					$5d^3$
		3	-2	$-\frac{1}{2}$	4
${}_{32}\text{Ge}$					
					$1s^2$
					$5f^{13}$
	4	0	0	$+\frac{1}{2}$	
	4	2	-1	$+\frac{1}{2}$	
					$6d^1$
${}_{70}\text{Yb}$					

5. What is true about the odd/evenness of the number of electrons in an atom and the resulting spin state of the last electron?

6. Demonstrate a working knowledge of how allowed values work by explaining why the p-block is 6 columns long and does not appear until  $n \geq 2$ . Make reference to **all four** quantum numbers. Use point form.
7. What would happen to the periodic table if an electron was able to spin four different ways (i.e.  $m_s = -3/2, -1/2, +1/2,$  and  $+3/2$ )? What would the widths of each "angular momentum" block become?



9. The Rydberg constant is itself a combination of different constants. Use the constants listed to determine the correct value of the Rydberg constant. Then perform a complete unit analysis. Be sure to start with the format "units ="

$$R = \frac{-e^4 m}{8\epsilon_0^2 h^3 c}$$

$e = 1.6022 \times 10^{-19}$  C (fundamental unit of charge)  
 $m = 9.110 \times 10^{-31}$  kg (resting mass of an electron)  
 $\pi = 3.1415926536$  (circumference / diameter for a circle)  
 $\epsilon_0 = 8.854 \times 10^{-12}$  C<sup>2</sup>N<sup>-1</sup>m<sup>-2</sup> (dielectric constant)  
 $h = 6.626 \times 10^{-34}$  Js (Planck's constant)  
 $c = 3.00 \times 10^8$  ms<sup>-1</sup> (speed of light)

$$J = \frac{\text{kgm}^2}{\text{s}^2}$$

$$N = \frac{\text{kgm}}{\text{s}^2}$$

10.

$$\frac{1}{\lambda} = 1.09737 \times 10^7 \text{ m}^{-1} \left[ \left( \frac{1}{n_i^2} \right) - \left( \frac{1}{n_f^2} \right) \right]$$

Look what light through yonder window duth break! Is it the sun? Is it the moon? No of course not, it is the fifth line in the Balmer Series of hydrogen. And what is the wavelength of yonder light? To figure that out, show a calculation for the appropriate initial and final states! Show a separate conversion to express your final answer in nanometers ( $1 \times 10^9 \text{ nm} = 1 \text{ m}$ ).

11. Using the equation from question 9, determine the initial and final states for:
- a) absorption of 379.70 nm
  - b) emission of 102.52 nm

wavelength	emitted or absorbed	$n_i$	$n_f$
379.70 nm	absorbed		
102.52 nm	emitted		