

## Acid Base Equilibrium Quantitative Problems

1. Calculate the  $[H_3O^{1+}]$ ,  $[OH^{1-}]$ , pH, and pOH of these solutions:
  - (a) 1.0 kmol/m<sup>3</sup> HCl
  - (b) 0.50 kmol/m<sup>3</sup> HNO<sub>3</sub>
  - (c) 0.0020 kmol/m<sup>3</sup> HClO<sub>4</sub>
  - (d) 1.5 X 10<sup>-4</sup> kmol/m<sup>3</sup> KOH
  - (e) a solution prepared by dissolving 0.040 g NaOH 2.0 L of solution
  - (f) a solution prepared by diluting 1.0 mL of 0.20 kmol/m<sup>3</sup> HCl to a total volume of 5.0 L
  - (g) a solution made by dissolving 0.10 mol Na<sub>2</sub>O in 1.0 L of solution
  
2. Calculate the  $[H_3O^{1+}]$ , pH, and percentage dissociation of these solutions:
  - (a) 1.0 kmol/m<sup>3</sup> HCN
  - (b) 0.001 kmol/m<sup>3</sup> HCN
  - (c) 1.0 kmol/m<sup>3</sup> HF
  - (d) 0.50 kmol/m<sup>3</sup> HNO<sub>2</sub>
  - (e) 0.5 kmol/m<sup>3</sup> HCHO<sub>2</sub>
  - (f) 0.50 kmol/m<sup>3</sup> H<sub>3</sub>BO<sub>3</sub>
  
3. A solution of hydrofluoric acid contains 2.0 g of HF per litre and has a pH of 2.2 What is the dissociation constant for HF?
  
4. A weak acid, HX, is a weak monoprotic acid. A 0.100 kmol/m<sup>3</sup> solution is 6.0% dissociated. What is the dissociation constant for the acid?
  
5. A 1.0 X 10<sup>-3</sup> kmol/m<sup>3</sup> solution of a weak acid, HX, is 20.0% dissociated.:
  - (a) What is the pH of the solution?
  - (b) What is the concentration of X<sup>1-</sup>?
  - (c) What is the dissociation constant for the acid?
  
6. Hypobromous acid (HBrO) has a dissociation constant of 2.0 X 10<sup>-9</sup>. A solution of HBrO has a pH of 4.8. What is the concentration of the solution?
  
7. Calculate the  $[OH^-]$ , pOH, and pH of these solutions:
  - (a) 1.0 kmol/m<sup>3</sup> NH<sub>3</sub>
  - (b) 0.10 kmol/m<sup>3</sup> aniline (C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub>),
  - (c) 5.0 X 10<sup>-2</sup> kmol/m<sup>3</sup> hydrazine (N<sub>2</sub>H<sub>4</sub>)
  - (d) 0.20 kmol/m<sup>3</sup> hydroxylamine (NH<sub>2</sub>OH)
  - (e) 1.5 kmol/m<sup>3</sup> trimethylamine ((CH<sub>3</sub>)<sub>3</sub>N)
  
8. Calculate the  $[OH^{1-}]$ , pOH and pH of these solutions:
  - (a) 0.10 kmol/m<sup>3</sup> Na<sub>2</sub>SO<sub>3</sub>
  - (b) 0.50 kmol/m<sup>3</sup> KCN
  - (c) 1.0 kmol/m<sup>3</sup> Na<sub>2</sub>CO<sub>3</sub>
  - (d) 0.05 kmol/m<sup>3</sup> NaC<sub>7</sub>H<sub>5</sub>O<sub>2</sub>
  - (e) 0.2 kmol/m<sup>3</sup> NaClO.
  
9. For this buffer solution:
  - (a) What is the pH of a solution made by combining 0.60 mol of acetic acid with 0.40 mol of sodium acetate in enough water to make one litre of solution?
  - (b) What is the pH of this solution if four additional litres of water is added to it?
  
10. What concentration of sodium acetate is required to prepare a solution in which the pH is 5.0 and the acetic acid is 0.10 kmol/m<sup>3</sup>.

11. What mass of  $\text{NH}_4\text{Cl}$  must be added to 0.500 L of 1.0 kmol/m<sup>3</sup>  $\text{NH}_3$  solution to yield a solution with a pH of 9.0? Assume no change in volume occurs.
12. A buffer solution is prepared by adding 1.0 mol of  $\text{NH}_4\text{Cl}$  to 1 litre of a solution containing 1.0 mol  $\text{NH}_3$ .
- (a) What is the pH of the solution?
  - (b) What is the pH of the solution resulting from the addition of 1.0 mmol of HCl to 10.0 mL of the buffer? Assume no volume change occurs.
  - (c) What is the pH of the solution resulting from the addition of 1.0 mmol NaOH to 1000 mL of the buffer? Assume no volume change.
  - (d) How many mL of 6 M HCl would be required to change the pH of one litre of buffer by 1 pH unit?